FDA/Johns Hopkins University/PhRMA Educational Workshop Drug Metabolizing Enzymes and Pharmacogennmic Testing-September 13-14, 2004 Rockville, MD

### Regulatory Issues in Genotyping Metabolizing Enzymes-CDER Perspective

Shiew-Mei Huang, Ph.D.

Deputy Office Director for Science

Office of Clinical Pharmacology & Biopharmaceutics, OPS

CDER, FDA

### Traditional Inefficiency in Drug Development

- Extremely high pre-IND failure rate
- Less than 1 in 5 IND's for NME's = NDA's
- Estimated cost per NME about \$800 million
- Time from IND to market around 8-10 years
- Multiple review cycles for most NME NDA's

Is this a systemic problem?
Is this the best that can be done?
What is needed?

esko, L, Pharmaceutical Sciences World Conference, Japan, May 18, 2004>

The product development problems we are seeing today can be addressed, ..... generation of ...... predictive tools.


Challenge and Opportunity on the Critical Path to New Medical Products

The new tools ..... such as bioinformatics, genomics, imaging technologies, and materials science.


http://www.fda.gov/oc/initiatives/criticalpath/whitepaper.html>


Clinical Pharmacology and Biopharmaceutics Review


**Extrinsic factors** Smoking/Diet **Environmental Drug-drug interaction** Medical Practice/ Regulatory **Intrinsic factors** Gender Genetics Race Organ Disease Age Dysfunction Pregnancy Lactation dapted from ICH E5, 1998: http://www.fda.gov/cder/guidance/2293fnl.pd





Shiew-Mei Huang, FDA/Johns Hopkins/PhRMA Workshop on "Drug Metabolizing Enzymes and Pharmacogenomics Testing, September 13, 2004, Rockville, MD















Shiew-Mei Huang, FDA/Johns Hopkins/PhRMA Workshop on "Drug Metabolizing Enzymes and Pharmacogenomics Testing, September 13, 2004, Rockville, MD


What <u>Pharmacogenomic Testing data</u> are needed when generating these initial PK data (or clinical data) that eventually support the labeling (or future labeling) for drugs metabolized by polymorphically distributed enzymes?



#### **Issues**

Are designations of EM and PM inadequate?
 do we need UM and IM?

#### CYP2D6



#### Issues (2)

- Are designations of EM and PM inadequate? do we need UM and IM?
- What data to support genotype-phenotype correlation?
- How could these designations be improved?
- What are the definitions of EM and PM for all enzymes?
  - "PM" applicable to all enzymes? CYP2C9?

15

#### Issues (3)

- What alleles need to be assessed before declaring a specific genotypes?
  - Can we extrapolate data from one ethnic/race group to the other?

#### CYP2C9

	Caucasian	Asian A	frican American	Hispanic
*2	8-19	0	3.2	12
*3	3.3-16	1.1-3.3	1.3	3.4
*5			0.2- <u>1.7</u>	0-0.5
*6			0.6- <u>1.5</u>	

#### Issues (4)

- •What information (PM, EM/others and specific alleles) is to be included in the drug label?
- To what extent should the test be commercially available?
- Why should the test be approved by the FDA to be in the labeling?

Guidance for Industry Pharmacogenomic Data Submissions

DRAFT GUIDANE

DRAFT GUIDANE


The plants of the plant of the plant


Known valid biomarker

.....CYP2D6 and TPMT (thiopurine methyltransferase)

: Guidance for Industry: Pharmacogenomic Data Submission:; >
http://www.fda.gov/cder/guidance/index.htm procedural (draft), November 2003; >

17





### **Case 1 (2)** Drug A

- · Based on PK and clinical data: the sponsor suggested no special dosing requirement for PM
- How critical is the accuracy in the PM and EM designation in evaluating these data?
- Alleles analyzed are limited and differ among studies
- Data analysis grouped based on EM and PM only
- Labeling mentions and makes recommendations based only EM and PM

### Case 2 Drug B (IND)

- Metabolized by CYP2C9
- the sponsor is conducting a drug interaction study with a CYP2C9 inhibitor, fluconazole
- Planning on retrospective CYP2C9 genotyping to identify impact on exposure of Drug B with fluconazole
- What alleles should be assessed in this study? Would the analysis of \*2,\*3,\*5 (reduced activity), or \*6 (null; but rare) appropriate?

# Case 3 Drug C (IND)

- Inhibits CYP2C19 and CYP3A
- the sponsor is conducting a drug interaction study with a CYP2C19/CYP3A substrate, nelfinavir
- Planning on retrospective CYP2C19 genotyping to identify impact on exposure of nelfinavir with drug C
- What alleles should be assessed in this study? Would the analysis of \*2 and \*3 appropriate?

Today's discussion on metabolic biomarkers

- · Known valid biomarkers
- CYP2D6, CYP2C9, CYP2C19
- · Probable valid biomarkers
- UGT1A1
- · Exploratory biomarkers
- ABCB1; CYP3A4/5
- Others

24

#### Special Considerations for Individual Metabolic Biomarkers Chair: S.-M. Huang, FDA

CYP2D6	D. Flockhart	Indiana Univ.
CYP2C9	P. Milos	Pfizer
CYP2C19	T. Andersson	AstraZeneca
UGT1A1	Mark Ratain	Univ. Chicago
CYP3A4/5	K. Thummel	Univ. Washington
P-gp and other transporters	D. Kroetz	UCSF
Haplotype Mapping of ADME Genes	D. Goldstein	UC, London

### **Discussion**

- In vitro in vivo correlation
- Genotype- phenotype correlation
- What alleles to measure?
- Haplotypes?

26

## Acknowledgement

OCPB Pharmacogenomics Working Group and OCPB reviewers (case examples)

A Bhattaram F Frueh S-M Huang M-J Kim I Lee L Lesko A Rahman K Roy A Rudman H Sun J Wei S Yasuda L Zhang S Nallani J Collins

**P** Hinderling

27